| WIDESPREAD SPECIES      | FREQUENCY                    | COMMENTS                         |
|-------------------------|------------------------------|----------------------------------|
| Canada Goose            | Abundant year-round resident |                                  |
| Mallard                 | Abundant year-round resident |                                  |
| Great Blue Heron        | Uncommon year-round resident |                                  |
| Sharp-shinned Hawk      | Uncommon year-round resident |                                  |
| Cooper's Hawk           | Uncommon year-round resident |                                  |
| Red-tailed Hawk         | Common year-round resident   |                                  |
| Rock Pigeon             | Abundant year-round resident |                                  |
| Mourning Dove           | Abundant year-round resident |                                  |
| Downy Woodpecker        | Common year-round resident   |                                  |
| Hairy Woodpecker        | Common year-round resident   |                                  |
| Blue Jay                | Common year-round resident   |                                  |
| American Crow           | Abundant year-round resident |                                  |
| Black-capped Chickadee  | Common year-round resident   |                                  |
| Tufted Titmouse         | Common year-round resident   |                                  |
| Red-breasted Nuthatch   | Uncommon year-round resident |                                  |
| White-breasted Nuthatch | Common year-round resident   |                                  |
| Eastern Bluebird        | Uncommon year-round resident |                                  |
| American Robin          | Common year-round resident   |                                  |
| European Starling       | Abundant year-round resident |                                  |
| Song Sparrow            | Common year-round resident   | Spring=Abundant; Winter=Uncommon |
| Northern Cardinal       | Common year-round resident   |                                  |
| House Finch             | Uncommon year-round resident |                                  |
| American Goldfinch      | Common year-round resident   |                                  |
| House Sparrow           | Abundant year-round resident |                                  |

#### **INDIVIDUAL SPECIES ACCOUNTS**

| Species Name | Occurrence | Approximate Arrival Date |
|--------------|------------|--------------------------|
|              |            | <pre></pre>              |

Habitat preference and additional information.

Likely location(s) to observe the species.

|            |                             | A 0/40 1 40/4=                    |
|------------|-----------------------------|-----------------------------------|
| Snow Goose | Seasonally abundant migrant | <b>↑</b> : 3/12; <b></b> ⊕: 10/15 |

More common in fall, feeding in harvested cornfields and resting on rivers, lakes and reservoirs. Often seen in large, flyover "skeins." Some overwinter where water remains open. Scan flocks for rare Ross' Goose.

Tomhannock Reservoir, Saratoga Lake, Fort Miller.

Brant Seasonally uncommon migrant ↑: mid-May; ↓: mid-Oct.

Passes through quickly. Majority of birds heard/seen as flyovers.

Cackling GooseSeasonally uncommon migrantSpring and Fall

Seen among migrating Canada Geese.

Mute Swan Locally uncommon year-round resident

Seen along Hudson River, occasionally on reservoirs, lakes, ponds, estuaries. Retreats southward in winter.

Tomhannock Reservoir, Stony Creek Reservoir, Basic Creek Reservoir.

Tundra Swan Rare winter visitant ↓: 11/1; ↑3/10

Rivers, lakes and reservoirs.

Saratoga Lake, Ft. Miller, Stillwater.

Wood Duck Locally common summer resident ↑: 3/10

Wooded swamps. Dabbler.

Vischer Ferry NHP, Mohawk-Hudson Bike-Hike Trail (Niskayuna), Black Creek Marsh.

Gadwall Seasonally uncommon migrant ↑: 3/10; ↓: 10/5

Shallow ponds, flooded fields. Dabbler.

Stanton Pond, Vischer Ferry NHP.

American Wigeon Seasonally uncommon migrant ↑: 3/10; ↓: 10/8

Shallow ponds, flooded fields. Dabbler.

Stanton Pond, Vischer Ferry NHP, Collins Lake.

Eurasian Wigeon Rare migrant

Seen in the company of American Wigeon.

American Black Duck Uncommon year-round resident

Shallow ponds, wetlands, marshes. Dabbler.

Black Creek Marsh, Vischer Ferry NHP, Stanton Pond

Blue-winged Teal Seasonally uncommon migrant ↑: 4/1; ↓: 8/20

Ponds and wetlands. Dabbler.

Black Creek Marsh, Vischer Ferry NHP, Cohoes Flats.

Northern Shoveler Seasonally uncommon migrant ↑: 3/15; ♦: 9/15

Ponds, wetlands, and flooded areas. Dabbler.

Stanton Pond, Vischer Ferry NHP, Collins Lake.

Northern Pintail Seasonally uncommon migrant ↑: 3/10; ↓: 10/1

Ponds and marshes. Dabbler.

Vischer Ferry NHP, Collins Lake, Stanton Pond.

Green-winged Teal Seasonally common migrant ↑: 3/10; ↓: 9/1

Ponds and marshes. Dabbler.

Vischer Ferry NHP, Collins Lake, Stanton Pond.

CanvasbackSeasonally rare migrant♠: 3/5; ♣: 11/5

Lakes and rivers. Mainly bypasses this area.

Ft. Miller, Saratoga Lake, Stillwater.

**Redhead** Seasonally rare migrant ↑: 3/5; ↓: 10/30

Lakes and rivers. Mainly bypasses this area.

Ft. Miller, Saratoga Lake, Stillwater.

**Ring-necked Duck** Seasonally common migrant ↑ 3/10; ↓: 10/10

Rivers, lakes, reservoirs, and ponds.

Vischer Ferry NHP, Stanton Pond, Collins Lake.

Greater Scaup Seasonally uncommon migrant ↑: 3/5; ♦: 10/20

Rivers, lakes and reservoirs.

Saratoga Lake, Tomhannock Reservoir, Ft. Miller.

**Lesser Scaup** Seasonally uncommon migrant ↑: 3/5; ♦: 10/20

Rivers, lakes and reservoirs.

Saratoga Lake, Vischer Ferry (Mohawk River), Ft. Miller.

Surf ScoterSeasonally rare migrant♣: 10/20

Rivers, lakes and reservoirs. Normally seen on fall migration only.

Mohawk River, Saratoga Lake, Tomhannock Reservoir.

White-winged ScoterSeasonally uncommon migrant↓: 10/20

Rivers, lakes and reservoirs. Normally seen on fall migration only.

Mohawk River, Saratoga Lake, Tomhannock Reservoir.

Rivers, lakes and reservoirs. Normally seen on fall migration only.

Mohawk River, Saratoga Lake, Tomhannock Reservoir.

**Long-tailed Duck** Seasonally uncommon migrant ↑: 3/15; ↓: 10/25

Rivers, lakes and reservoirs.

Basic Creek Reservoir, Myosotis Lake, Mohawk River.

**Bufflehead** Seasonally uncommon migrant ↑: 3/10; ↓: 11/1

Rivers, lakes and reservoirs.

Basic Creek Reservoir, Vischer Ferry (Mohawk River), Ft. Miller.

Common GoldeneyeCommon winter resident↓: 11/5

Rivers, lakes and reservoirs. Scan for rare Barrow's Goldeneye.

**Red-breasted Merganser** Seasonally uncommon migrant ↑: 3/10; ♦: 10/20

Very uncommon, usually 1-4 birds, April/November.

Mohawk River, Tomhannock Reservoir, Saratoga Lake.

Hooded Merganser | Locally uncommon year-round resident | migrants↑: 3/15; ⊕: 10/10

Virtually any body of water. Most of the population breeds north of our area, and winters south of the region. A few overwinter if water remains open.

Common MerganserCommon winter resident↓: 10/10

Virtually any body of water, October-April.

**Ruddy Duck** Seasonally uncommon migrant ↑: 3/10; ↓: 10/10

More common in fall.

Stanton Pond, Tomhannock Reservoir, Basic Creek Reservoir.

Ruffed Grouse Locally uncommon year-round resident

Early succession mixed deciduous forests, with small clearings.

Partridge Run WMA, Saratoga NHP, Thacher Park

Wild Turkey Locally uncommon year-round resident

Hardwood forests with scattered openings.

Black Creek Marsh (and vicinity), Albany Pine Bush Preserve, Saratoga NHP

| Red-throated Loon | Seasonally rare migrant | <b>↑: 4/6;                                   </b> |
|-------------------|-------------------------|---|
|-------------------|-------------------------|---|

Rivers, lakes and reservoirs.

Tomhannock Reservoir, Collins Lake, Mohawk River.

| Common Loon | Seasonally uncommon migrant | <b>↑</b> : 3/25; <b>↓</b> : 10/10 |
|-------------|-----------------------------|-----------------------------------|
|-------------|-----------------------------|-----------------------------------|

Rivers, lakes and reservoirs.

Alcove Reservoir, Saratoga Lake, Mohawk River.

| Pied-billed Grebe Seasonally uncommon migrant ↑: 3/14; ↓: 10/1 |
|--|
|--|

Marshes, shallow ponds.

Collins Lake, Ann Lee Pond, Vischer Ferry NHP.

| Horned Grebe Seasonally uncommon migrant ↑: 3/10; ↓: 10/10 |  |
|--|--|
|--|--|

Rivers, lakes and reservoirs.

Collins Lake, Mohawk River, Tomhannock Reservoir.

| Red-necked Grebe | Seasonally uncommon migrant | <b>↑: 3/25;                                    </b> |
|------------------|-----------------------------|---|
|------------------|-----------------------------|---|

Rivers, lakes and reservoirs.

Tomhannock Reservoir, Hudson River.

| <b>Double-crested Cormorant</b> | Common summer resident | <b>↑: 3/24</b> |
|---------------------------------|------------------------|----------------|
|---------------------------------|------------------------|----------------|

Widespread on reservoirs, lakes and rivers.

| <b>Great Cormorant</b> | Uncommon early spring visitant | Late-Feb. through March |
|------------------------|--------------------------------|-------------------------|

On the Hudson River from about Coeymans southward. Favors river channel markers as perch. A few fall records, mostly juveniles.

Embought Bay (Catskill), Germantown.

| American Bittern Locally uncommon so | nmer resident 1: 4/15 |
|--------------------------------------|-----------------------|
|--------------------------------------|-----------------------|

In large cattail marshes.

Black Creek Marsh, Cline Rd. marsh (Oppenheim).

| Least Bittern | Locally uncommon summer resident | <b>↑: 5/7</b> |
|---------------|----------------------------------|---------------|
|---------------|----------------------------------|---------------|

In large cattail marshes.

Black Creek Marsh, Vischer Ferry NHP.

| <b>Great Egret</b> | Locally uncommon summer visitant | <b>↑: 4/5; ←&gt;: 7/20</b> |
|--------------------|----------------------------------|----------------------------|

Small numbers seen during spring migration; larger numbers during post-breeding dispersal. Any shallow water along rivers, ponds, lakes.

Vischer Ferry NHP, Mohawk River.

| Snowy Egret  | Rare summer visitant | ⇔Late July- September |
|--|----------------------|-----------------------|
| Along Inour the Hudson and Mahauk Biyars, Cohoos Flats |                      |                       |

Along/near the Hudson and Mohawk Rivers, Cohoes Flats.

Green Heron Locally common summer resident ↑: 4/20

Shallow ponds, marshes, wooded wetlands.

Black Creek Marsh, Vischer Ferry NHP, Five Rivers EEC.

Black-crowned Night-Heron Seasonally rare migrant ↑: 4/19; ↓: 8/7

Rivers, lakes, wooded marshes.

Vischer Ferry NHP, Cohoes Flats, Black Creek Marsh.

Black Vulture Uncommon year-round resident

Increasing due to range expansion. Retreats southward in winter, though many are now overwintering as far north as Albany County.

Thacher Park, Five Rivers EEC.

 Turkey Vulture
 Common summer resident
 ↑: 2/12

Widespread in small numbers. Retreats southward in winter.

OspreySeasonally uncommon summer resident↑: 4/1

Rivers, lakes and ponds.

Hudson and Mohawk Rivers.

Bald Eagle Locally uncommon year-round resident

Numbers increasing. Rivers, lakes and ponds.

Hudson and Mohawk Rivers, Tomhannock Reservoir.

Golden Eagle Seasonally rare migrant ↑: March; ↓: late October

Main breeding population migrates west of the area. Quebec & Labrador population crosses our area en route to/from wintering grounds.

Mississippi Kite Locally rare summer resident ↑: present by end of May

Range expanding. Breeding pair established in Montgomery County, NY in 2009/2010. Also present in southeast NH and northern CT.

| Northern Harrier | Locally uncommon year-round resident |  |
|------------------|--------------------------------------|--|
|------------------|--------------------------------------|--|

Open habitat, especially grasslands and marshes.

Ft. Edward Grasslands, Coxsackie Grasslands, Five Rivers EEC.

| Northern Goshawk | Locally rare year-round resident |  |
|------------------|----------------------------------|--|
| Hortine Goshawk  | Locally rate year round resident |  |

Very localized breeder throughout the region. Prefers dense, contiguous forested areas.

Dyken Pond EEC, Partridge Run WMA.

| Red-shouldered Hawk | Locally uncommon summer resident | ↑: 3/12 |
|---------------------|----------------------------------|---------|
|                     |                                  |         |

Deciduous woodlands, often near rivers and swamps. Most birds retreat just south of the area during the winter months, a few try to overwinter. Red-shouldered Hawks return to the same nesting territory year after year.

Vischer Ferry NHP, Stanton Pond, Partridge Run WMA

| <b>Broad-winged Hawk</b> | Locally uncommon summer resident | ↑: 4/20;   migrants: 9/10 |
|--------------------------|----------------------------------|---------------------------|
|                          |                                  |                           |

Breeds in forested areas, especially at higher elevations. Migrates in large groups. Can be seen in great numbers at fall hawk watches.

Western Albany County, Rensselaer Plateau.

| Rough-legged Hawk | Locally uncommon winter resident |  |
|-------------------|----------------------------------|--|
| Open grasslands.  |                                  |  |

Coxsackie, Ft. Edward, Northumberland.

| American Kestrel | Locally uncommon year-round resident |  |
|------------------|--------------------------------------|--|
|------------------|--------------------------------------|--|

Open habitat, with scattered trees for surveying prey. Often seen on utility wires.

Saratoga NHP, Coxsackie Grasslands, Ft. Edward.

| Merlin | Locally uncommon year-round resident |  |
|--------|--------------------------------------|--|
|        |                                      |  |

Along rivers. Also seen at hawk watches during migration. Increasingly found in more populated areas.

Cohoes Flats/Peebles Island, Coxsackie.

|--|

Along bodies of water, primarily the Hudson River. Nests in boxes on bridges, naturally on cliff faces.

Cohoes, Albany, Thacher Park.

| Virginia Rail | Locally uncommon summer resident | <b>1</b> : 4/10 |
|---------------|----------------------------------|-----------------|
| B             |                                  |                 |

Cattail marshes. Calls alert observers to their presence, though they are frequently seen through diligent observation, and during post-breeding forays.

Black Creek Marsh, Vischer Ferry NHP.

| Sora | Locally uncommon summer resident | <b>1</b> : 4/14 |
|------|----------------------------------|-----------------|
|------|----------------------------------|-----------------|

Cattail marshes. More often heard than seen.

Black Creek Marsh, Vischer Ferry NHP, RamsHorn-Livingston Sanctuary.

| Common Gallinule | Locally uncommon summer resident | <b>↑: 4/20</b> |
|------------------|----------------------------------|----------------|
|                  |                                  |                |

Cattail marshes.

Black Creek Marsh, Vischer Ferry NHP, Cline Rd. marsh (Oppenheim).

| American Coot    | Seasonally uncommon migrant | <b>↑</b> : 3/10; <b>↓</b> : 10/20 |
|------------------|-----------------------------|-----------------------------------|
| Lakes and ponds. |                             |                                   |

Vischer Ferry NHP, Collins Lake, Stanton Pond.

| Black-bellied Plover Seasonally rare migrant | <b>↑</b> : 5/15; <b>↓</b> : 8/15 |
|--|----------------------------------|
|--|----------------------------------|

Mudflats, flooded fields.

Mohawk River, Wright's Loop.

| American Golden-Plover | Seasonally rare migrant | <b>.</b> |
|------------------------|-------------------------|----------|
|------------------------|-------------------------|----------|

Most likely during fall migration. Pastures, rocky flats.

Cohoes Flats, Myosotis Lake.

| Semipalmated Plover       | Seasonally uncommon migrant | <b>↑</b> : 5/14; <b></b> ⊕: 7/20 |
|---------------------------|-----------------------------|----------------------------------|
| Mudflats, flooded fields. |                             |                                  |

Cohoes, Vischer Ferry (Mohawk River), Staats Island Rd. (Schodack).

| Killdeer   | Locally common summer resident | <b>↑: 2/25</b> |
|--|--------------------------------|----------------|
| NAC de consecutivo de contense e contense e contense e contense de contense de contense de contense e contense |                                |                |

Widespread in shortgrass and gravel areas, rocky flats.

| Spotted Sandpiper | Locally common summer resident | <b>1</b> : 4/19 |  |
|-------------------|--------------------------------|-----------------|--|
|-------------------|--------------------------------|-----------------|--|

Ponds, creeks, shorelines. Usually a singleton.

Vischer Ferry NHP, Peebles Island, Five Rivers EEC.

| Solitary Sandpiper | Seasonally common migrant | <b>↑</b> : 4/27; <b></b> ⊕: 7/15 |  |
|--------------------|---------------------------|----------------------------------|--|
|--------------------|---------------------------|----------------------------------|--|

Shallow, muddy, vegetated ponds and creeks, flooded areas.

Vischer Ferry NHP, Five Rivers EEC, Staats Island Rd. (Schodack).

Any shallow water habitat.

Vischer Ferry NHP, Cohoes Flats, Stanton Pond.

| Lesser Yellowlegs | Seasonally uncommon migrant | <b>↑</b> : 4/22; <b>♦</b> : 7/15 |
|-------------------|-----------------------------|----------------------------------|
|-------------------|-----------------------------|----------------------------------|

Any shallow water habitat.

Vischer Ferry NHP, Cohoes Flats, Papscanee Island North.

Short grass fields and pastures, especially in agricultural areas.

Ft. Edward Grasslands, Fulton & Montgomery County farms/pastures. Notable breeders in West Ames, Montgomery County.

| Sanderling | Seasonally rare migrant | <b>.</b> 8/15 |
|------------|-------------------------|---------------|
|------------|-------------------------|---------------|

More common during fall migration. At water's edge or on rocky flats.

Mohawk, Hudson Rivers.

| Semipalmated Sandpiper   Seasonally uncommon migrant | <b>↓: 7/15</b> |
|--|----------------|
|--|----------------|

More common during fall migration. Mudflats, any shallow water habitat.

Cohoes Flats, Coxsackie Grasslands, Mohawk River.

| Least Sandpiper | Seasonally uncommon migrant | <b>↑: 5/1;                                   </b> |
|-----------------|-----------------------------|---|
|-----------------|-----------------------------|---|

Mudflats, ponds, any shallow water habitat.

Cohoes Flats, Vischer Ferry NHP, Stanton Pond.

| White-rumped Sandpiper | Seasonally rare migrant | <b>↓</b> : 8/1 |
|------------------------|-------------------------|----------------|
|------------------------|-------------------------|----------------|

More common during fall migration. Mudflats, ponds, any shallow water habitat.

Cohoes Flats, Vischer Ferry NHP, Stanton Pond.

| Baird's Sandpiper Seasonally rare migrant | <b>.</b> 8/20 |
|---|---------------|
|---|---------------|

More common during fall migration. Mudflats, grassy marshes, and dry grassy areas near lakes and ponds.

Myosotis Lake, Cohoes Flats, Coxsackie Grasslands.

More common during fall migration. Prefers weedy margins of lakes and ponds, along with muddy agricultural fields. Prolonged fall migration. Adults from mid-July through August; juveniles in September and October.

Myosotis Lake, Cohoes Flats, Stanton Pond.

DunlinSeasonally rare migrant↑: 4/27; ♦: 10/1

Mudflats, flooded areas.

Staats Island Rd., Cohoes Flats, Mohawk River.

Stilt SandpiperSeasonally rare migrant♣: 9/1

More common during fall migration. Mudflats, flooded fields, shallow ponds and pools, and marshes.

Cohoes Flats, The Crossings (Colonie), Myosotis Lake.

More common during fall migration. Mud flats and flooded agricultural fields.

Cohoes Flats, Vischer Ferry NHP, Myosotis Lake.

Wilson's Snipe Uncommon summer resident ↑: 4/1

Wet, grassy areas, edges of shallow ponds.

Vischer Ferry NHP, Five Rivers EEC, Black Creek Marsh.

 American Woodcock
 Uncommon summer resident
 ↑: 3/8

Along wooded edge habitat.

Albany Pine Bush, Vischer Ferry NHP, Five Rivers EEC.

Bonaparte's Gull Seasonally uncommon migrant ↑: 4/5; ↓: 10/10

Rivers and lakes. A few wanderers occasionally seen in summer.

Cohoes Flats, Saratoga Lake, Mohawk River.

Ring-billed Gull Abundant year-round resident

Rivers, lakes, landfills, parking lots. Summer residents are non-breeding individuals. Ring-billed Gulls breed in the Great Lakes region.

Herring Gull Abundant year-round resident

Rivers, lakes, landfills, parking lots. Summer residents are non-breeding individuals. Herring Gulls breed on Long Island and in the Adirondacks/Lake Champlain Region.

Iceland GullRare winter visitantNov.- March

Occasionally appears among winter gulls along Hudson/Mohawk Rivers.

Lesser Black-backed GullRare winter visitantNov.- March

Occasionally appears among winter gulls along Hudson/Mohawk Rivers.

Glaucous Gull Rare winter visitant Nov.- March

Occasionally appears among winter gulls along Hudson/Mohawk Rivers.

Great Black-backed Gull Locally uncommon year-round resident

Rivers, reservoirs, lakes. Summer residents are non-breeding individuals. Great Black-backed Gulls breed along the Atlantic Coast from Long Island to the Canadian Maritime Provinces.

Caspian Tern Rare spring migrant/summer visitant ↑: 4/10; ← : 7/10

Lakes and rivers. A few wanderers occasionally seen in mid-late summer.

Black Tern Rare spring migrant ↑: 5/15

Lakes and rivers.

Common Tern Rare spring migrant ↑: 5/1

Lakes and rivers.

Saratoga Lake, Mohawk River.

Yellow-billed Cuckoo Uncommon summer resident ↑: 5/13

Open woodlands with clearings and dense scrubby vegetation, often along water.

Schodack Island SP, Vischer Ferry NHP, Black Creek Marsh.

Black-billed Cuckoo Uncommon summer resident ↑: 5/9

Deciduous woodlands and thickets.

Saratoga NHP, Five Rivers EEC.

Eastern Screech-Owl Uncommon year-round resident

Any wooded habitat with tree cavities, or nesting boxes. Often suns itself in those openings during the day.

Black Creek Marsh, Vosburgh Marsh, Five Rivers EEC.

Great Horned Owl Uncommon year-round resident

Open and secondary-growth woodlands and agricultural areas.

Five Rivers EEC, Vischer Ferry NHP, Saratoga NHP.

Snowy Owl Rare winter visitant

Irruptive species moves southward (typically immature birds), especially in years with high breeding success.

Barred Owl Locally uncommon year-round resident

Mixed woodland habitat with large trees.

Saratoga NHP, Five Rivers EEC, Partridge Run WMA.

Locally rare year-round resident

Open forest adjacent to grasslands/shrubland.

Open country grasslands, marshes.

Ft. Edward Grasslands, Coxsackie Grasslands

Forests, densely vegetated habitats.

Western Albany County.

| Common Nighthawk | Seasonally uncommon migrant | <b>↑</b> : 5/22; <b>♦</b> : 8/10 |
|------------------|-----------------------------|----------------------------------|
| - 11 1           |                             |                                  |

Declining. Formerly a resident breeder, now most commonly seen in migratory flight towards sundown.

| Eastern Whip-poor-will Locally rare summer resident | <b>1</b> : 4/29 |  |
|---|-----------------|--|
|---|-----------------|--|

Open, mature deciduous or mixed forest.

Stanton Pond, Malta Tech Park.

| Chimney Swift | Locally common summer resident | <b>1</b> : 4/27 |
|---------------|--------------------------------|-----------------|
|---------------|--------------------------------|-----------------|

Commonly nests in urban areas, forages over open areas.

Cohoes Flats, Five Rivers EEC.

| <b>Ruby-throated Hummingbird</b> | Uncommon summer resident | <b>1</b> : 4/30 |
|----------------------------------|--------------------------|-----------------|
|----------------------------------|--------------------------|-----------------|

Forest edges, old fields, orchards, meadows and backyards.

Five Rivers EEC, Schodack Island SP, Vischer Ferry NHP.

| Belted Kingfisher | Locally uncommon year-round resident |  |
|-------------------|--------------------------------------|--|

Streams, rivers, ponds, and lakes. Overwinters as long as open water is available.

Five Rivers EEC, Vischer Ferry NHP, Cohoes Flats.

| Red-headed Woodpecker | Locally rare year-round resident |  |
|-----------------------|----------------------------------|--|
|-----------------------|----------------------------------|--|

Open oak or pine savannah, wooded beaver swamps.

Albany Pine Bush Preserve

| Red-bellied Woodpecker | Common year-round resident |  |
|------------------------|----------------------------|--|
|------------------------|----------------------------|--|

Forests, woodlands, and backyards.

Five Rivers EEC, Vischer Ferry NHP, Collins Lake.

Young forests and edge habitat. A few birds linger through winter.

Five Rivers EEC, Saratoga NHP, Thacher Park.

Northern Flicker Common year-round resident

Woodlands, forest edges, open areas with scattered trees. Some birds regularly overwinter.

Five Rivers EEC, Saratoga NHP, Albany Pine Bush Preserve.

Pileated Woodpecker Uncommon year-round resident

Mature woodlands, suburban woodlots with large trees.

Thacher Park, Five Rivers EEC, Vischer Ferry NHP.

Olive-sided FlycatcherSeasonally rare migrant↑: 5/15; ↓: 8/20

Forest edges and openings, ponds, with tall snags.

Five Rivers EEC, Albany Pine Bush Preserve, Ann Lee Pond.

Eastern Wood-Pewee Locally uncommon summer resident ↑: 5/12

All types of woodlands.

Thacher Park, Black Creek Marsh, Partridge Run WMA.

Yellow-bellied Flycatcher | Locally rare summer resident | ↑: 5/20; ♣ migrants: 8/20

Bogs, swamps, coniferous forest. Breeds at the highest elevations of the Catskill Mountains.

Plateau Mountain, Hunter Mountain.

Alder FlycatcherLocally uncommon summer resident↑: 5/15

Wet thickets, especially of alder and birch.

Five Rivers EEC, Black Creek Marsh, Saratoga NHP.

Willow Flycatcher Locally uncommon summer resident ↑: 5/16

Moist, shrubby areas often with standing water.

Five Rivers EEC, Black Creek Marsh, Vischer Ferry NHP.

 Least Flycatcher
 Locally uncommon summer resident
 ↑: 5/7

Semi-open woodlands, shrubby fields.

Papscanee Island NP, Five Rivers EEC, Vischer Ferry NHP.

Eastern Phoebe Common summer resident ↑: 3/20

Open woodlands, backyards, parks especially near water.

Five Rivers EEC, Vischer Ferry NHP, Mohawk-Hudson Bike/Hike Trail.

Open deciduous woodlands, riparian corridors, wooded swamps, urban areas with large trees.

Schodack Island SP, Five Rivers EEC, Vischer Ferry NHP.

Eastern Kingbird Locally uncommon summer resident 1: 4/25

Fields, pastures, grasslands, and marshes with shrubs. Usually near water.

Saratoga NHP, Black Creek Marsh, Vischer Ferry NHP.

Northern Shrike Locally uncommon winter resident ↓: 11/1

Open country with tall perches, shrubby fields, marshes.

Western Albany County, Coxsackie Grasslands, Black Creek Marsh.

Yellow-throated Vireo Locally uncommon summer resident ↑: 4/27

Edge habitat in mature deciduous woods, usually near water.

Schodack Island SP, Black Creek Marsh, Vischer Ferry NHP.

Blue-headed Vireo Locally uncommon summer resident ↑: 4/18

Cool, higher elevation forests. Common Spring/Fall migrant.

Dyken Pond EEC, Cherry Plain SP, Thacher Park.

Warbling Vireo Locally uncommon summer resident ↑: 4/26

Mixed-deciduous woodlands along streams, ponds, and marshes.

Vischer Ferry NHP, Black Creek Marsh, Schodack Island SP.

Philadelphia Vireo Seasonally uncommon migrant ↑: 5/10; ↓: 9/1

Deciduous woodlands. More often seen on fall migration.

Vischer Ferry NHP, Five Rivers EEC.

Red-eyed VireoLocally common summer resident↑: 5/6

Deciduous and mixed-deciduous forests, urban areas with large trees.

Hannacroix Creek Preserve, Five Rivers EEC, Albany Pine Bush Preserve.

Fish Crow Locally uncommon year-round resident

Along rivers, lakes, and landfills.

Hudson River, Peebles Island SP, The Crossings (Colonie).

Common Raven Locally uncommon year-round resident

A variety of habitats, mostly outside of urban areas.

Thacher Park, Five Rivers EEC, Tomhannock Reservoir.

Horned Lark Locally uncommon year-round resident

Open, barren country; airports, plowed agricultural fields, roadside edges in winter.

Coxsackie Flats, Saratoga County Airport, Northumberland farmlands (north of Bacon Hill).

| Purple Martin | Locally rare summer resident | <b>↑</b> : 4/20 |
|---------------|------------------------------|-----------------|

Colonial breeders in nest boxes and martin houses, especially in open areas near water.

Saratoga Lake and vicinity.

Open areas near water and fields, especially wooded swamps and shorelines.

Five Rivers EEC, Black Creek Marsh, Cohoes Flats.

| Northern Rough-winged | Locally uncommon summer resident | <b>1</b> : 4/7 |
|-----------------------|----------------------------------|----------------|
| Swallow               |                                  |                |

Open habitats with nesting surfaces such as river banks and human structures.

Mohawk River Lock 7, Cohoes Flats, Tomhannock Reservoir.

| Bank Swallow | Locally uncommon summer resident | <b>1</b> : 4/18 |
|--------------|----------------------------------|-----------------|

River banks or sand/gravel quarries.

Coxsackie Grasslands, Papscanee Island NP, Peebles Island SP.

| Cliff Swallow | Locally uncommon summer resident | <b>↑: 5/1</b> |
|---------------|----------------------------------|---------------|
|---------------|----------------------------------|---------------|

Habitats with open foraging areas, especially near bridges and other man-made structures for nesting.

Alcove Reservoir, Schoharie Creek bridges.

| Barn Swallow | Locally common summer resident | <b>↑: 4/7</b> |
|--------------|--------------------------------|---------------|
|--------------|--------------------------------|---------------|

Open habitat including parks, agricultural areas, lakes, and ponds, with nearby structures for nesting.

Black Creek Marsh, Stanton Pond, The Crossings (Colonie).

Evergreen or mixed evergreen-deciduous forests.

Five Rivers EEC, Thacher Park, Albany Pine Bush Preserve.

| Carolina Wren | Locally uncommon year-round resident |  |
|---------------|--------------------------------------|--|

Dense vegetation in wooded areas, primarily at lower elevations.

Five Rivers EEC, Vischer Ferry NHP, Peebles Island SP.

| House Wren Locally uncommon summer resident ↑: 4/19 |
|---|
|---|

Edge habitat with trees, shrubs, and tangles; also farms and backyards.

Five Rivers EEC, Black Creek Marsh, Vischer Ferry NHP.

Winter Wren Locally uncommon summer resident ↑: 4/10

Riparian corridors, coniferous forests with fallen trees, logs. A few birds overwinter; most winter just to our south, return north in April and May at lower elevations first, then expand northward and to higher elevations in June.

Five Rivers EEC, Vischer Ferry NHP, Thacher Park.

Sedge WrenLocally rare summer resident↑: mid-May

Damp, tall-grass meadows with scattered bushes.

Ft. Edward grasslands, Sharon Springs (and vicinity).

Marsh WrenLocally uncommon summer resident↑: 4/13

Cattail marshes. A few birds may attempt to overwinter.

Black Creek Marsh, Vischer Ferry NHP, Mohawk-Hudson Bike/Hike Trail (Niskayuna).

Golden-crowned Kinglet Locally uncommon year-round resident

Breeds in higher elevation spruce forest, more often seen Sept. through April when retreating to lower elevation mixed woodlands.

Partridge Run WMA, Thacher Park, Five Rivers EEC.

 Ruby-crowned Kinglet
 Seasonally uncommon migrant
 ↑: 4/7

Breeds mainly to our north, in spruce-fir forests; a few breeding reports from the northern Catskills. Winters just to the south of our area, occasional Jan.-Mar. reports along the Hudson River.

Five Rivers EEC, Black Creek Marsh, Vischer Ferry NHP.

Deciduous wooded habitats, especially near water.

Black Creek Marsh, Vischer Ferry NHP, Schodack Island SP.

 Veery
 Locally uncommon summer resident
 ↑: 5/4

Damp, deciduous forests and riparian corridors. Along with the Wood Thrush, lives at the lowest elevations of the resident thrushes.

Holt Preserve, Schodack Island SP, Papscanee Island NP.

Gray-cheeked Thrush Seasonally rare migrant ↑: 5/12; ↓: 9/25

More often reported on fall migration, usually heard overhead in nocturnal flight.

**Bicknell's Thrush** Locally rare summer resident ↑: 5/24

Breeds at the highest elevations of the Catskill Mountains.

Plateau Mountain, Hunter Mountain.

| Swainson's Thrush                    | Locally uncommon summer resident           | <b>↑</b> : 5/4; <b>↓</b> : 9/5 |
|--------------------------------------|--|--------------------------------|
| 3 VV a i i i i i a 3 i i i i a 3 i i | Locally afficultifier satisfier restaction | · J/ ¬, % · J/ J               |

Breeds in spruce-fir forests in the Catskill Mountains, higher than the Hermit Thrush but below Bicknell's. More commonly observed on fall migration in lower elevation woodlands.

Partridge Run WMA, Hunter Mountain, Plateau Mountain.

Coniferous and deciduous forests with open understory for foraging. Hermit Thrushes occupy the middle elevations of the resident thrushes. Some birds attempt to overwinter.

Thacher Park, Dyken Pond EEC, Partridge Run WMA.

| Wood Thrush | Locally uncommon summer resident | <b>1</b> : 4/27 |
|-------------|----------------------------------|-----------------|
|-------------|----------------------------------|-----------------|

Mature forests with an open floor and moist soil, with water nearby.

Black Creek Marsh, Schodack Island SP, Papscanee Island NP.

Dense shrubs, vine tangles, and thickets of young trees.

Five Rivers EEC, Schodack Island SP, Vischer Ferry NHP.

| ear-round resident |
|--------------------|
|--------------------|

Open ground with shrubby vegetation such as hedges, fruiting bushes, and thickets.

The Crossings (Colonie), Five Rivers EEC, Saratoga NHP

| Brown Thrasher | Uncommon summer resident | <b>↑: 4/10</b> |
|----------------|--------------------------|----------------|
|----------------|--------------------------|----------------|

Thickets, hedgerows, and forest edges.

Five Rivers EEC, Black Creek Marsh, Rensselaer Tech Park.

| American Pipit | Seasonally uncommon migrant | <b>.</b> ⊕: 9/20 |  |
|----------------|-----------------------------|------------------|--|
|----------------|-----------------------------|------------------|--|

More often seen on fall migration. Recently plowed fields, mudflats, and river courses.

Stanton Pond, Myosotis Lake, Staats Island Rd.

| Bohemian Waxwing | Rare winter visitant |  |
|------------------|----------------------|--|
|------------------|----------------------|--|

Irruptive species feeds on mountain ash, crabapples and other fruit-bearing trees. Sometimes occurs in mixed company with Cedar Waxwing. Usually seen in northern portions of our area.

| Cedar Waxwing | Common year-round resident |  |
|---------------|----------------------------|--|
|---------------|----------------------------|--|

Deciduous, coniferous, and mixed woodlands, particularly areas along streams. Diet primarily consists of fruits, but is seen catching insects over ponds and streams in summer.

| Lapland Longspur | Locally rare winter resident | ⊕: Dec early March |
|------------------|------------------------------|--------------------|
|------------------|------------------------------|--------------------|

Plowed fields, stubble, and open grasslands. Often associates with Horned Lark.

Northumberland farm fields, Schaghticoke farm fields, Coxsackie Flats.

Open weedy and grassy fields, and along shores of lakes.

Ft. Edward Flats, Northumberland farm fields, Coxsackie Flats.

| Blue-winged Warbler | Locally uncommon summer resident | <b>↑: 4/26</b> |
|---------------------|----------------------------------|----------------|

Early to mid-succession habitats, especially abandoned farmland, shrublands, and forest clearings.

Five Rivers EEC, Saratoga NHP, Hannacroix Creek Preserve.

| Golden-winged Warbler | Locally rare summer resident | <b>↑:5/7</b> |
|-----------------------|------------------------------|--------------|
|-----------------------|------------------------------|--------------|

Nearly extirpated from the region as a breeding species. Early successional habitats such as recently abandoned farms, shrublands, and clearcuts.

Northeastern Washington County near VT border.

| Tennessee Warbler Seasonally uncommon | nigrant ↑: 5/10; ↓: 9/1 |
|---------------------------------------|-------------------------|
|---------------------------------------|-------------------------|

Open woodlands. More often seen on fall migration.

Thacher Park, Partridge Run WMA, Five Rivers EEC.

| Orange-crowned Warbler | Seasonally rare migrant | <b>.</b> ⊕: 9/24 |
|------------------------|-------------------------|------------------|
|------------------------|-------------------------|------------------|

Thickets and dense tangles of shrubs and vines. Nearly all sightings are on fall migration, in late Sept-Oct. for this late migrant.

Five Rivers EEC, Peebles Island SP, Papscanee Island NP.

| Nashville Warbler | Locally uncommon summer resident | ↑: 4/25;   migrants: 9/1 |
|-------------------|----------------------------------|--------------------------|

Second-growth deciduous or mixed forest with shrubby undergrowth. Usually seen on migration, a few breeding locations have been found at higher elevations in Rensselaer, Albany, Greene and Schoharie Counties.

Grafton Lakes SP, Burnt-Rossman Hills SF, Cherry Plain SP (and vicinity).

| Northern Parula | Seasonally uncommon migrant | <b>↑</b> : 4/29; <b>⊎</b> : 8/25 |  |
|-----------------|-----------------------------|----------------------------------|--|
|-----------------|-----------------------------|----------------------------------|--|

Mature forests along streams, swamps, and other bottomlands. Normally seen on migration only, but breeds both south and north of our area. A few June/July reports from the Schoharie Valley.

Max Shaul SP (and vicinity).

Yellow Warbler Locally common summer resident ↑: 4/23

Shrubby thickets and woods, particularly along watercourses and in wetlands, especially where willows are present.

Vischer Ferry NHP, Black Creek Marsh, Papscanee Island NP.

 Chestnut-sided Warbler
 Locally uncommon summer resident
 ↑: 4/30

Early successional deciduous woods and scrubby forests.

Partridge Run WMA, Albany Pine Bush Preserve, Schodack Island SP.

Magnolia WarblerLocally uncommon summer resident↑: 5/3; ♣ migrants: 8/25

Usually seen on migration. Breeds in small conifers, especially young spruces, in purely coniferous stands or mixed forest. A few breeders at higher elevations on the Rensselaer Plateau, western Albany County, and Schoharie County.

Dyken Pond EEC, Burnt-Rossman Hills SF, Partridge Run WMA.

Cape May WarblerSeasonally uncommon migrant↑: 5/7; ↓: 8/25

Coniferous forests and mixed woodlands, but nearly always around spruce trees.

Five Rivers, Partridge Run WMA, Camp Pinnacle (and vicinity).

Black-throated Blue Warbler Locally uncommon summer resident ↑: 5/1; ♣ migrants: 8/25

Usually seen on migration. Breeds in mature deciduous and mixed coniferous-deciduous woodlands with a thick understory, usually at higher elevations.

Partridge Run WMA, Burnt-Rossman Hills SF, Dyken Pond EEC.

Yellow-rumped Warbler Locally uncommon summer resident ↑: 4/12; 

 migrants: 9/1

Mature coniferous and mixed coniferous-deciduous woodlands. Very common during spring and fall migration, a few breeders are found at higher elevations on the Rensselaer Plateau, Northern Saratoga County, western Albany County, and the Schoharie Valley.

Dyken Pond EEC, Cherry Plain SP, Burnt-Rossman Hills SF.

Black-throated Green Warbler | Locally uncommon summer resident | ↑: 4/23: ↓ migrants: 8/25 | Coniferous forests and transitional coniferous-deciduous forest, usually at higher elevations..

Dyken Pond EEC, Cherry Plain SP, Deer Mountain Nature Trail.

**Blackburnian Warbler** Locally uncommon summer resident ↑: 4/28; ♣ migrants: 8/25

Mature coniferous and mixed coniferous/deciduous forests at higher elevations.

Dyken Pond EEC, Cherry Plain SP, Burnt-Rossman Hills SF.

| Pine Warbler | Locally uncommon summer resident | <b>↑: 4/4</b> |
|--------------|----------------------------------|---------------|
|--------------|----------------------------------|---------------|

Pine or mixed pine-deciduous forest.

Albany Pine Bush Preserve, Five Rivers EEC, Wilton Wildlife Preserve.

| Prairie Warbler | Locally uncommon summer resident | <b>1</b> : 4/27 |
|-----------------|----------------------------------|-----------------|
|-----------------|----------------------------------|-----------------|

Shrubby habitats and open fields, often with pine trees present.

Five Rivers EEC, Albany Pine Bush Preserve, Black Creek Marsh (and vicinity).

| Palm Warbler | Seasonally uncommon migrant | <b>↑: 4/7;                                   </b> |  |
|--------------|-----------------------------|---|--|
|--------------|-----------------------------|---|--|

Second growth and thicket habitats, and open fields.

Five Rivers EEC, Black Creek Marsh, Vischer Ferry Preserve.

| Bay-breasted Warbler | Seasonally rare migrant | <b>↑</b> : 5/11; <b>↓</b> : 9/1 |
|----------------------|-------------------------|---------------------------------|
|----------------------|-------------------------|---------------------------------|

Coniferous woodlands, more often seen on fall migration.

Five Rivers EEC, Thacher Park, Albany Pine Bush Preserve.

| Diackpoil warpier   Seasonally uncommon migrant   1 1. 5/12 9/10 | Blackpoll Warbler | Seasonally uncommon migrant | <b>↑: 5/12;                                    </b> |
|--|-------------------|-----------------------------|---|
|--|-------------------|-----------------------------|---|

Various forest, woodland, scrub and brushy habitats. A small population breeds at higher elevations in the Catskill Mountains.

Five Rivers EEC, Albany Pine Bush Preserve, Vischer Ferry NHP.

| Cerulean Warbler | Locally uncommon summer resident | <b>↑: 5/5</b> |
|------------------|----------------------------------|---------------|
|------------------|----------------------------------|---------------|

Forests with tall deciduous trees and open understory.

Max Shaul SP (and vicinity), Schodack Island SP, Hannacroix Creek Preserve.

| Black-and-white Warbler | Locally uncommon summer resident | <b>↑: 4/27</b> |
|-------------------------|----------------------------------|----------------|
|-------------------------|----------------------------------|----------------|

Mature and second-growth deciduous and mixed forests.

Schodack Island SP, Holt Preserve, Max Shaul SP (and vicinity).

| American Redstart | Locally common summer resident | <b>1</b> : 4/27 |
|-------------------|--------------------------------|-----------------|
|-------------------|--------------------------------|-----------------|

Moist second growth deciduous forest, with abundant shrubs.

Schodack Island SP, Papscanee Island NP, Vischer Ferry NHP.

| Worm-eating Warbler | Locally uncommon summer resident | <b>↑: 5/3</b> |
|---------------------|----------------------------------|---------------|
|                     |                                  |               |

Steep slopes with dense understory.

Deer Mountain Nature Trail, Holt Preserve, Bennett Hill Preserve.

| Ovenbird | Locally common summer resident | <b>1</b> : 4/26 |
|----------|--------------------------------|-----------------|

Mature deciduous and mixed deciduous and coniferous forests.

Dyken Pond EEC, Schodack Island SP, Five Rivers.

| Northern Waterthrush | Locally uncommon summer resident | <b>↑: 4/26</b> |
|----------------------|----------------------------------|----------------|
|----------------------|----------------------------------|----------------|

Thickets near slow-moving streams, ponds, swamps, and bogs.

Vischer Ferry NHP, Partridge Run WMA, Grafton Lakes SP.

| Louisiana Waterthrush | Locally uncommon summer resident | <b>↑: 4/9</b> |
|-----------------------|----------------------------------|---------------|
|-----------------------|----------------------------------|---------------|

Gravel-bottomed streams flowing through hilly, deciduous forest.

Five Rivers EEC, Hannacroix Creek Preserve, Thacher Park.

| <b>Connecticut Warbler</b> | Seasonally rare migrant | <b>.</b> 9/7 |
|----------------------------|-------------------------|--------------|
|----------------------------|-------------------------|--------------|

Scrub and thicket habitats. All sightings are on fall migration.

Vischer Ferry NHP, Albany Pine Bush Preserve, Dyken Pond EEC.

| Mourning Warbler | Locally uncommon summer resident | ↑: 5/16;   migrants: 9/1 |
|------------------|----------------------------------|--------------------------|
|------------------|----------------------------------|--------------------------|

Second-growth forested areas, with moderately closed canopy and thick vegetation.

Partridge Run WMA, Cherry Plain SP, Berlin Mountain.

| Common Yellowthroat | Locally common summer resident | <b>↑</b> : 4/25 |
|---------------------|--------------------------------|-----------------|
|---------------------|--------------------------------|-----------------|

Common in thick vegetation from wetlands to pine forests. Frequently near water.

Five Rivers EEC, Albany Pine Bush Preserve, Vischer Ferry NHP.

|                |                                  | A = 15         |
|----------------|----------------------------------|----------------|
| Hooded Warbler | Locally uncommon summer resident | <b>↑</b> : 5/6 |

Hardwood forests with some shrub understory.

Holt Preserve, Bennett Hill Preserve, Partridge Run WMA.

| Wilson's Warbler | Seasonally uncommon migrant | <b>↑</b> : 5/7; <b>↓</b> : 8/26 |
|------------------|-----------------------------|---------------------------------|

Secondary growth and scrub thickets, pond edges and riparian habitats.

Schodack Island SP, Five Rivers EEC, Vischer Ferry NHP.

|  | Canada Warbler | Locally uncommon summer resident | ↑: 5/5;   migrants: 8/15 |
|--|----------------|----------------------------------|--------------------------|
|--|----------------|----------------------------------|--------------------------|

Found in moist thickets in riparian corridors, brushy ravines, forest bogs.

Max Shaul SP (and vicinity), Cherry Plain SP, Partridge Run WMA.

| Eastern Towhee | Locally uncommon summer resident | <b>1</b> : 4/7 |
|----------------|----------------------------------|----------------|
|----------------|----------------------------------|----------------|

Forest edges, overgrown fields and woodlands, and scrubby backyards or thickets. Winters just to the south of our area.

Five Rivers EEC, Albany Pine Bush Preserve, Saratoga NHP.

| American Tree Sparrow Locally uncommon winter resident | American Tree Sparrow | Locally uncommon winter resident | <b>₺</b> : 10/23 |
|--|-----------------------|----------------------------------|------------------|
|--|-----------------------|----------------------------------|------------------|

Weedy fields with hedgerows or shrubs, along forest edges, or near marshes.

Five Rivers EEC, Vischer Ferry NHP, Saratoga NHP.

| <b>Chipping Sparrow</b> | Locally common summer resident | <b>↑: 4/10</b> |
|-------------------------|--------------------------------|----------------|
|-------------------------|--------------------------------|----------------|

Open woodlands, forests with grassy clearings, parks.

Albany Pine Bush Preserve, The Crossings (Colonie), Saratoga NHP.

| Clay-colored Sparrow | Locally rare summer resident | <b>↑: 5/2</b> |
|----------------------|------------------------------|---------------|
|----------------------|------------------------------|---------------|

Brushy and weedy, dry habitats such as pastures and field edges. Population gradually expanding eastward from the Great Lakes region.

Five Rivers EEC, Sharon Springs (and vicinity), Ft. Edward Grasslands.

| Field Coorney. | Locally was a name on a was as a said and | <b>↑.</b> 2/20 |
|----------------|---|----------------|
| Field Sparrow  | Locally uncommon summer resident          | " ": 3/20      |

Brushy pastures and old fields. Winters just to the south of our area.

Five Rivers EEC, Albany Pine Bush Preserve, Saratoga NHP.

| Vesper Sparrow Locally rare summer resident 1:4 | 4/13 |
|---|------|
|---|------|

Open habitats with grass.

Ft. Edward Grasslands, Saratoga County Airport, Coxsackie Grasslands.

| Savannah Sparrow | Locally uncommon summer resident | <b>↑: 4/11</b> |
|------------------|----------------------------------|----------------|
|                  |                                  |                |

Open areas with low vegetation, grasslands with few trees. A few birds overwinter, primarily in southern parts of our area.

Staats Island Rd. (Schodack), The Crossings (Colonie), Saratoga County Airport.

|                     |                                  | 1              |
|---------------------|----------------------------------|----------------|
| Grasshopper Sparrow | Locally uncommon summer resident | <b>↑</b> : 5/5 |

Open grasslands with patches of bare ground.

Saratoga County Airport, Ft. Edward Grasslands.

| Fox Sparrow Seas | sonally uncommon migrant | <b>↑:</b> 3/14; <b>↓:</b> 10/14 |
|------------------|--------------------------|---------------------------------|
|------------------|--------------------------|---------------------------------|

Thickets of dense, shrubby growth. Winters just to the south of our area.

Five Rivers EEC, Ann Lee Pond, Vischer Ferry NHP.

| Lincoln's Sparrow | Seasonally uncommon migrant | <b>↑</b> : 5/7; <b>↓</b> : 9/3 |
|-------------------|-----------------------------|--------------------------------|

Brushy areas, thickets, hedgerows, clearings, and scrubby areas. More common on fall migration.

Five Rivers EEC, Vischer Ferry NHP, Coxsackie Grasslands.

| Swamp Sparrow   Locally uncommon summer resident   1.4/2 |
|--|
|--|

Various wetlands, including marshes, bogs, and swamps. Winters just to the south of our area.

Black Creek Marsh, Vischer Ferry NHP, Bog Meadow Brook Nature Trail.

White-throated Sparrow Locally uncommon year-round resident

Forest edges, thickets, overgrown fields, and parks. Retreats to higher elevations in summer.

Dyken Pond EEC, Cherry Plain SP, Partridge Run WMA.

White-crowned Sparrow Seasonally uncommon migrant ↑: 4/28; ♦: 9/27

Thickets, weedy fields, and agricultural fields. A few birds overwinter, primarily in southern parts of our area.

Five Rivers EEC, Vischer Ferry NHP.

Dark-eyed Junco Locally uncommon year-round resident

Coniferous forests, open woodlands, fields and parks. Common in low elevations in winter.

Dyken Pond EEC, Grafton Lakes SP, Burnt-Rossman Hills SF.

 Scarlet Tanager
 Locally uncommon summer resident
 ↑: 5/3

Deciduous and mixed deciduous/coniferous woodlands.

Five Rivers EEC, Vischer Ferry NHP, Albany Pine Bush Preserve.

Rose-breasted Grosbeak Locally uncommon summer resident 1: 4/27

Deciduous and mixed woodlands, especially at the edges.

Five Rivers EEC, Schodack Island SP, Vischer Ferry NHP.

 Indigo Bunting
 Locally uncommon summer resident
 ↑: 5/9

Brushy and weedy areas, on the edges of woods and fields, and along powerline cuts.

Albany Pine Bush Preserve, Malta Tech Park, Saratoga NHP.

**Bobolink** Locally uncommon summer resident ↑: 4/29

Open grasslands and hay fields.

Five Rivers EEC, Saratoga NHP, Coxsackie Grasslands.

| Red-winged Blackbird | Locally abundant summer resident | Mid-Feb. through mid-Dec. |
|----------------------|----------------------------------|---------------------------|
|----------------------|----------------------------------|---------------------------|

Marshes, agricultural fields, pastures, and wet grasslands. Retreats slightly south of the area for a short time.

Areawide.

| Eastern Meadowlark | Locally uncommon summer resident | <b>1</b> : 3/26 |
|--------------------|----------------------------------|-----------------|
|--------------------|----------------------------------|-----------------|

Farm fields, grasslands, and wet fields. A few birds attempt to overwinter.

Coxsackie Flats, Saratoga NHP, Ft. Edward Grasslands.

Wet areas, including flooded woods, swamps, marshes and the edges of ponds.

Vischer Ferry NHP, Ann Lee Pond, Bog Meadow Brook Nature Trail.

| Common Grackle | Locally abundant summer resident | Late Feb. through mid-Nov. |
|----------------|----------------------------------|----------------------------|
|                |                                  |                            |

Agricultural fields, parks, woodland edges, marshes, and suburban lawns. Retreats slightly south of the area for a short time.

Areawide.

| Brown-headed Cowbird   | Locally uncommon summer resident  | Mid-Feb. through mid-Nov.          |
|------------------------|-----------------------------------|------------------------------------|
| DIOWII IICAACA COWDIIA | Locally allcommon sammer restacht | I wild I CD. till ough illia ivov. |

Woodland edges, brushy thickets, fields, pastures, orchards, and residential areas. A few birds overwinter, mainly in southern parts of our area.

Areawide.

| Orchard Oriole | Locally uncommon summer resident | <b>1</b> : 5/6 |
|----------------|----------------------------------|----------------|
|----------------|----------------------------------|----------------|

Deciduous trees along streams, rivers and lakes, and near gardens and orchards.

Five Rivers EEC, Normanskill Farm, Saratoga NHP.

| <b>Baltimore Oriole</b> | Locally uncommon summer resident | <b>↑: 4/27</b> |
|-------------------------|----------------------------------|----------------|
|                         |                                  |                |

Open woodland, forest edge, orchards, and stands of trees along rivers, in parks, and in suburban areas.

Five Rivers EEC, Schodack Island SP, Vischer Ferry NHP.

| Pine Grosbeak Rare winter visitant ↓: Nov Feb. |
|--|
|--|

Irruptive species feeds on fruit-bearing trees. Often found in urban/suburban areas with ornamental fruit tree plantings.

| Purple Finch | Locally uncommon year-round resident |  |
|--------------|--------------------------------------|--|

Moist, cool evergreen forests, and mixed forests, along wooded streams.

Partridge Run WMA, Cherry Plain SP, Burnt-Rossman Hills SF.

**Red Crossbill** Locally rare year-round resident

Mature coniferous forests. Wanders widely between years to find a good cone crop. Will breed throughout most of the year, whenever it finds areas with an abundance of cones.

Burnt-Rossman Hills SF, Partridge Run WMA.

White-winged Crossbill Rare winter visitant ↓: Nov.- March

Coniferous forests, especially where large crops of spruce and tamarack cones can be found. Wanders widely between years to find a good cone crop. Will breed throughout most of the year, whenever it finds areas with an abundance of cones. Breeding reports from higher elevations of Rensselaer County.

Irruptive species. Winter range extremely variable as birds seek unpredictable food sources. Open woodlands, scrubby and weedy fields, especially with birch trees.

Pine Siskin Locally uncommon year-round resident

Coniferous or mixed coniferous forest. Irrupts southward and to lower elevations in search of seeds. Opportunistic and adaptable, foraging in weedy fields, scrubby thickets, or suburban yards and gardens. A few breeding locations in the higher elevations of western Albany and eastern Rensselaer Counties.

Irruptive species. Winters in coniferous or deciduous forests, and in urban and suburban areas.

#### Citations:

eBird. 2009. eBird: An online database of bird distribution and abundance [web application]. Version 2. eBird, Ithaca, New York. Available: <a href="http://www.ebird.org">http://www.ebird.org</a>. (Accessed: November 2012).

Cornell Laboratory of Ornithology and National Audubon Society. 2009. eBird: An online database of bird distribution and abundance. Avian Knowledge Network. Ithaca, NY. <a href="www.avianknowledge.net">www.avianknowledge.net</a>. Data accessed: November 2012.

New York State Breeding Bird Atlas 2000 [Internet]. 2000 - 2005. Release 1.0. Albany (New York): New York State Department of Environmental Conservation. [updated 2007 Jun 11; cited 2012 Nov 29]. Available from: <a href="http://www.dec.ny.gov/animals/7312.html">http://www.dec.ny.gov/animals/7312.html</a>.

Cornell Laboratory of Ornithology. All About Birds: Bird Guide: Life Histories. <a href="http://www.allaboutbirds.org/Page.aspx?pid=1189">http://www.allaboutbirds.org/Page.aspx?pid=1189</a>.

Hudson-Mohawk Bird Club. <a href="http://hmbc.net">http://hmbc.net</a> & HMBirds Yahoo Group. Anecdotal sightings. <a href="http://groups.yahoo.com/group/hmbird">http://groups.yahoo.com/group/hmbird</a>.

#### Appendix A: List of Birds by Approximate Arrival Date

| JANUARY- all month       |                                      |  |
|--------------------------|--------------------------------------|--|
| Tundra Swan              | Rare winter visitant                 |  |
| American Black Duck      | Uncommon year-round resident         |  |
| Common Goldeneye         | Common winter resident               |  |
| Common Merganser         | Common winter resident               |  |
| Ruffed Grouse            | Locally uncommon year-round resident |  |
| Wild Turkey              | Locally uncommon year-round resident |  |
| Black Vulture            | Locally uncommon year-round resident |  |
| Bald Eagle               | Locally uncommon year-round resident |  |
| Northern Harrier         | Locally uncommon year-round resident |  |
| Northern Goshawk         | Locally rare year-round resident     |  |
| Rough-legged Hawk        | Locally uncommon winter resident     |  |
| American Kestrel         | Locally uncommon year-round resident |  |
| Merlin                   | Locally uncommon year-round resident |  |
| Ring-billed Gull         | Abundant year-round resident         |  |
| Herring Gull             | Abundant year-round resident         |  |
| Peregrine Falcon         | Locally uncommon year-round resident |  |
| Iceland Gull             | Rare winter visitant                 |  |
| Lesser Black-backed Gull | Rare winter visitant                 |  |
| Glaucous Gull            | Rare winter visitant                 |  |
| Great Black-backed Gull  | Locally uncommon year-round resident |  |
| Eastern Screech-Owl      | Uncommon year-round resident         |  |

| Great Horned Owl       | Uncommon year-round resident         |
|------------------------|--------------------------------------|
| Snowy Owl              | Rare winter visitant                 |
| Barred Owl             | Locally uncommon year-round resident |
| Long-eared Owl         | Locally rare year-round resident     |
| Short-eared Owl        | Locally uncommon winter resident     |
| Northern Saw-whet Owl  | Locally rare year-round resident     |
| Belted Kingfisher      | Locally uncommon year-round resident |
| Red-headed Woodpecker  | Locally rare year-round resident     |
| Red-bellied Woodpecker | Common year-round resident           |
| Northern Flicker       | Locally uncommon year-round resident |
| Pileated Woodpecker    | Uncommon year-round resident         |
| Northern Shrike        | Locally uncommon winter resident     |
| Fish Crow              | Locally uncommon year-round resident |
| Common Raven           | Locally uncommon year-round resident |
| Horned Lark            | Locally uncommon year-round resident |
| Brown Creeper          | Locally uncommon year-round resident |
| Carolina Wren          | Locally uncommon year-round resident |
| Golden-crowned Kinglet | Locally uncommon year-round resident |
| Northern Mockingbird   | Locally uncommon year-round resident |
| Bohemian Waxwing       | Rare winter visitant                 |
| Cedar Waxwing          | Common year-round resident           |
| Lapland Longspur       | Rare winter resident                 |
| Snow Bunting           | Locally uncommon winter resident     |
| American Tree Sparrow  | Locally uncommon winter resident     |

| White-throated Sparrow | Locally uncommon year-round resident |  |
|------------------------|--------------------------------------|--|
| Dark-eyed Junco        | Locally uncommon year-round resident |  |
| Pine Grosbeak          | Rare winter visitant                 |  |
| Purple Finch           | Locally uncommon year-round resident |  |
| Red Crossbill          | Locally rare year-round resident     |  |
| White-winged Crossbill | Rare winter visitant                 |  |
| Common Redpoll         | Locally uncommon winter visitant     |  |
| Pine Siskin            | Locally uncommon year-round resident |  |
| Evening Grosbeak       | Locally uncommon winter visitant     |  |
| FEBRUARY               |                                      |  |
| Turkey Vulture         | 2/12                                 |  |
| Red-winged Blackbird   | 2/15                                 |  |
| Brown-headed Cowbird   | 2/20                                 |  |
| Great Cormorant        | 2/24                                 |  |
| Common Grackle         | 2/25                                 |  |
| Killdeer               | 2/25                                 |  |
| MARCH                  |                                      |  |
| Mute Swan              | 3/1                                  |  |
| Canvasback             | 3/5                                  |  |
| Redhead                | 3/5                                  |  |
| Greater Scaup          | 3/5                                  |  |
| Lesser Scaup           | 3/5                                  |  |
| American Woodcock      | 3/8                                  |  |
| Wood Duck              | 3/10                                 |  |

| American Wigeon          | 3/10 |  |
|--------------------------|------|--|
| Eurasian Wigeon          | 3/10 |  |
| Northern Pintail         | 3/10 |  |
| Green-winged Teal        | 3/10 |  |
| Bufflehead               | 3/10 |  |
| Ring-necked Duck         | 3/10 |  |
| Red-breasted Merganser   | 3/10 |  |
| Ruddy Duck               | 3/10 |  |
| Horned Grebe             | 3/10 |  |
| American Coot            | 3/10 |  |
| Snow Goose               | 3/12 |  |
| Red-shouldered Hawk      | 3/12 |  |
| Tree Swallow             | 3/12 |  |
| Gadwall                  | 3/14 |  |
| Pied-billed Grebe        | 3/14 |  |
| Fox Sparrow              | 3/14 |  |
| Northern Shoveler        | 3/15 |  |
| Hooded Merganser         | 3/15 |  |
| Long-tailed Duck         | 3/15 |  |
| Golden Eagle             | 3/15 |  |
| Eastern Phoebe           | 3/20 |  |
| Field Sparrow            | 3/20 |  |
| Rusty Blackbird          | 3/20 |  |
| Double-crested Cormorant | 3/24 |  |

| Common Loon                   | 3/25 |  |
|-------------------------------|------|--|
| Red-necked Grebe              | 3/25 |  |
| Eastern Meadowlark            | 3/26 |  |
| APRIL                         |      |  |
| Blue-winged Teal              | 4/1  |  |
| Osprey                        | 4/1  |  |
| Wilson's Snipe                | 4/1  |  |
| Yellow-bellied Sapsucker      | 4/1  |  |
| Swamp Sparrow                 | 4/2  |  |
| Pine Warbler                  | 4/4  |  |
| Great Egret                   | 4/5  |  |
| Bonaparte's Gull              | 4/5  |  |
| Red-throated Loon             | 4/6  |  |
| Hermit Thrush                 | 4/6  |  |
| Northern Rough-winged Swallow | 4/7  |  |
| Barn Swallow                  | 4/7  |  |
| Ruby-crowned Kinglet          | 4/7  |  |
| Palm Warbler                  | 4/7  |  |
| Eastern Towhee                | 4/7  |  |
| Louisiana Waterthrush         | 4/9  |  |
| Virginia Rail                 | 4/10 |  |
| Greater Yellowlegs            | 4/10 |  |
| Winter Wren                   | 4/10 |  |
| Brown Thrasher                | 4/10 |  |

| Chipping Sparrow             | 4/10 |  |
|------------------------------|------|--|
| Caspian Tern                 | 4/10 |  |
| Savannah Sparrow             | 4/11 |  |
| Yellow-rumped Warbler        | 4/12 |  |
| Marsh Wren                   | 4/13 |  |
| Vesper Sparrow               | 4/13 |  |
| Sora                         | 4/14 |  |
| American Bittern             | 4/15 |  |
| Blue-headed Vireo            | 4/18 |  |
| Blue-gray Gnatcatcher        | 4/18 |  |
| Bank Swallow                 | 4/18 |  |
| Black-crowned Night-Heron    | 4/19 |  |
| Spotted Sandpiper            | 4/19 |  |
| House Wren                   | 4/19 |  |
| Green Heron                  | 4/20 |  |
| Common Gallinule             | 4/20 |  |
| Broad-winged Hawk            | 4/20 |  |
| Purple Martin                | 4/20 |  |
| Upland Sandpiper             | 4/21 |  |
| Lesser Yellowlegs            | 4/22 |  |
| Yellow Warbler               | 4/23 |  |
| Black-throated Green Warbler | 4/23 |  |
| Eastern Kingbird             | 4/25 |  |
| Nashville Warbler            | 4/25 |  |

| Common Yellowthroat       | 4/25 |  |
|---------------------------|------|--|
| Warbling Vireo            | 4/26 |  |
| Gray Catbird              | 4/26 |  |
| Blue-winged Warbler       | 4/26 |  |
| Ovenbird                  | 4/26 |  |
| Northern Waterthrush      | 4/26 |  |
| Solitary Sandpiper        | 4/27 |  |
| Dunlin                    | 4/27 |  |
| Chimney Swift             | 4/27 |  |
| Yellow-throated Vireo     | 4/27 |  |
| Wood Thrush               | 4/27 |  |
| Prairie Warbler           | 4/27 |  |
| Black-and-white Warbler   | 4/27 |  |
| American Redstart         | 4/27 |  |
| Rose-breasted Grosbeak    | 4/27 |  |
| Baltimore Oriole          | 4/27 |  |
| Blackburnian Warbler      | 4/28 |  |
| White-crowned Sparrow     | 4/28 |  |
| Eastern Whip-poor-will    | 4/29 |  |
| Northern Parula           | 4/29 |  |
| Bobolink                  | 4/29 |  |
| Ruby-throated Hummingbird | 4/30 |  |
| Chestnut-sided Warbler    | 4/30 |  |
| MAY                       |      |  |

| Least Sandpiper             | 5/1     |  |
|-----------------------------|---------|--|
| Great-crested Flycatcher    | 5/1     |  |
| Cliff Swallow               | 5/1     |  |
| Black-throated Blue Warbler | 5/1     |  |
| Clay-colored Sparrow        | 5/2 (?) |  |
| Magnolia Warbler            | 5/3     |  |
| Worm-eating Warbler         | 5/3     |  |
| Scarlet Tanager             | 5/3     |  |
| Veery                       | 5/4     |  |
| Swainson's Thrush           | 5/4     |  |
| Cerulean Warbler            | 5/5     |  |
| Canada Warbler              | 5/5     |  |
| Grasshopper Sparrow         | 5/5     |  |
| Red-eyed Vireo              | 5/6     |  |
| Hooded Warbler              | 5/6     |  |
| Orchard Oriole              | 5/6     |  |
| Least Bittern               | 5/7     |  |
| Least Flycatcher            | 5/7     |  |
| Golden-winged Warbler       | 5/7 (?) |  |
| Cape May Warbler            | 5/7     |  |
| Wilson's Warbler            | 5/7     |  |
| Lincoln's Sparrow           | 5/7     |  |
| Black-billed Cuckoo         | 5/9     |  |
| Indigo Bunting              | 5/9     |  |

| Brant                     | 5/10     |  |
|---------------------------|----------|--|
| Common Tern               | 5/10     |  |
|                           |          |  |
| Philadelphia Vireo        | 5/10     |  |
| Tennessee Warbler         | 5/10     |  |
| Bay-breasted Warbler      | 5/11     |  |
| Eastern Wood-Pewee        | 5/12     |  |
| Gray-cheeked Thrush       | 5/12     |  |
| Blackpoll Warbler         | 5/12     |  |
| Yellow-billed Cuckoo      | 5/13     |  |
| Semipalmated Plover       | 5/14     |  |
| Black Tern                | 5/15     |  |
| Alder Flycatcher          | 5/15     |  |
| Mississippi Kite          | 5/15 (?) |  |
| Black-bellied Plover      | 5/15     |  |
| Olive-sided Flycatcher    | 5/15     |  |
| Sedge Wren                | 5/15 (?) |  |
| Willow Flycatcher         | 5/16     |  |
| Mourning Warbler          | 5/16     |  |
| Yellow-bellied Flycatcher | 5/20     |  |
| Common Nighthawk          | 5/22     |  |
| Bicknell's Thrush         | 5/24     |  |
| JUNE                      |          |  |
| JULY                      |          |  |
| Least Sandpiper           | 7/10     |  |

| Caspian Tern              | 7/10 |  |
|---------------------------|------|--|
| Semipalmated Sandpiper    | 7/15 |  |
| Lesser Yellowlegs         | 7/15 |  |
| Solitary Sandpiper        | 7/15 |  |
| Short-billed Dowitcher    | 7/15 |  |
| Semipalmated Plover       | 7/20 |  |
| Great Egret               | 7/20 |  |
| Snowy Egret               | 7/22 |  |
| AUGUST                    |      |  |
| Pectoral Sandpiper        | 8/1  |  |
| White-rumped Sandpiper    | 8/1  |  |
| Greater Yellowlegs        | 8/5  |  |
| Black-crowned Night-Heron | 8/7  |  |
| Common Nighthawk          | 8/10 |  |
| Sanderling                | 8/15 |  |
| Black-bellied Plover      | 8/15 |  |
| Canada Warbler            | 8/15 |  |
| Blue-winged Teal          | 8/20 |  |
| Baird's Sandpiper         | 8/20 |  |
| Olive-sided Flycatcher    | 8/20 |  |
| Yellow-bellied Flycatcher | 8/20 |  |
| Northern Parula           | 8/25 |  |
| Magnolia Warbler          | 8/25 |  |
| Cape May Warbler          | 8/25 |  |

| Black-throated Blue Warbler  | 8/25 |  |
|------------------------------|------|--|
| Black-throated Green Warbler | 8/25 |  |
| Blackburnian Warbler         | 8/25 |  |
| Wilson's Warbler             | 8/26 |  |
| SEPTEMBER                    |      |  |
| Green-winged Teal            | 9/1  |  |
| Philadelphia Vireo           | 9/1  |  |
| Tennessee Warbler            | 9/1  |  |
| Nashville Warbler            | 9/1  |  |
| Yellow-rumped Warbler        | 9/1  |  |
| Bay-breasted Warbler         | 9/1  |  |
| Mourning Warbler             | 9/1  |  |
| American Golden-Plover       | 9/1  |  |
| Stilt Sandpiper              | 9/1  |  |
| Lincoln's Sparrow            | 9/3  |  |
| Swainson's Thrush            | 9/5  |  |
| Connecticut Warbler          | 9/7  |  |
| Blackpoll Warbler            | 9/10 |  |
| Palm Warbler                 | 9/10 |  |
| Northern Shoveler            | 9/15 |  |
| American Pipit               | 9/20 |  |
| Orange-crowned Warbler       | 9/24 |  |
| Gray-cheeked Thrush          | 9/25 |  |
| Rusty Blackbird              | 9/25 |  |

| White-crowned Sparrow  | 9/27  |  |
|------------------------|-------|--|
| OCTOBER                |       |  |
| Pied-billed Grebe      | 10/1  |  |
| Dunlin                 | 10/1  |  |
| Northern Pintail       | 10/1  |  |
| Gadwall                | 10/5  |  |
| American Wigeon        | 10/8  |  |
| Horned Grebe           | 10/10 |  |
| Brant                  | 10/10 |  |
| Red-throated Loon      | 10/10 |  |
| Ring-necked Duck       | 10/10 |  |
| Common Merganser       | 10/10 |  |
| Hooded Merganser       | 10/10 |  |
| Ruddy Duck             | 10/10 |  |
| Bonaparte's Gull       | 10/10 |  |
| Evening Grosbeak       | 10/10 |  |
| Common Loon            | 10/10 |  |
| Fox Sparrow            | 10/14 |  |
| Snow Goose             | 10/15 |  |
| Red-necked Grebe       | 10/15 |  |
| American Coot          | 10/20 |  |
| Red-breasted Merganser | 10/20 |  |
| Greater Scaup          | 10/20 |  |
| Lesser Scaup           | 10/20 |  |

| Surf Scoter              | 10/20 |  |
|--------------------------|-------|--|
| White-winged Scoter      | 10/20 |  |
| Black Scoter             | 10/20 |  |
| American Tree Sparrow    | 10/23 |  |
| Long-tailed Duck         | 10/25 |  |
| Golden Eagle             | 10/25 |  |
| Redhead                  | 10/30 |  |
| NOVEMBER                 |       |  |
| Bufflehead               | 11/1  |  |
| Northern Shrike          | 11/1  |  |
| Rough-legged Hawk        | 11/1  |  |
| Tundra Swan              | 11/1  |  |
| Snow Bunting             | 11/1  |  |
| Common Redpoll           | 11/1  |  |
| Short-eared Owl          | 11/1  |  |
| White-winged Crossbill   | 11/1  |  |
| Canvasback               | 11/5  |  |
| Common Goldeneye         | 11/5  |  |
| Iceland Gull             | 11/10 |  |
| Lesser Black-backed Gull | 11/10 |  |
| Glaucous Gull            | 11/10 |  |
| Pine Grosbeak            | 11/15 |  |
| Lapland Longspur         | 11/30 |  |